

## COMPUTING A NEW TOPOLOGICAL INDEX OF NANO STRUCTURES

MODJTABA GHORBANI, MARYAM JALALI\*

*Department of Mathematics, Faculty of Science, Shahid Rajaei Teacher Training University, Tehran, 16785 – 136, I. R. Iran*

*Institute of Nanoscience and Nanotechnology, University of Kashan, Kashan 87317-51167, I. R. Iran*

Let  $G$  be a graph and  $e = uv$  be an edge of  $G$ . The GA index of  $G$  is defined as

$$GA(G) = \sum_{e \in E} \frac{2\sqrt{du dv}}{du + dv}. \text{ In this paper we compute the GA index of } TUC_4C_8(S) \text{ nanotube.}$$

(Received September 11, 2009; accepted September 30, 2009)

*Keywords:* Topological Indices, nanotube, nanotori, GA Index.

### 1. Introduction

A nanostructure is an object of intermediate size between molecular and microscopic structures. It is a product derived through engineering at the molecular scale. The most important of these new materials are carbon nanotubes<sup>1-3</sup>. They have remarkable electronic properties and many other unique characteristics. For these reasons it is of interest to study the mathematical properties of these materials.

Mathematical chemistry is a branch of theoretical chemistry for discussion and prediction of the molecular structure using mathematical methods without necessarily referring to quantum mechanics. Chemical graph theory is a branch of mathematical chemistry which applies graph theory to mathematical modeling of chemical phenomena<sup>4-7</sup>. This theory had an important effect on the development of the chemical sciences.

A topological index is a numeric quantity from the structural graph of a molecule. Usage of topological indices in chemistry began in 1947 when chemist Harold Wiener developed the most widely known topological descriptor, the Wiener index, and used it to

determine physical properties of types of alkanes known as paraffin<sup>8</sup>. The Hosoya polynomial of a graph  $G$  is defined as  $W(G;x) = \sum_{u,v \in V(G)} x^{d(u,v)}$ , where  $d(u,v)$  denotes the length of a minimum path between  $u$  and  $v$ . In the paper [9] Hosoya used the name Wiener polynomial while some authors later used the name Hosoya polynomial.

Let  $G$  be a graph and  $e = uv$  be an edge of  $G$ . The GA index of  $G$  was introduced by D. Vukičević and co-authors<sup>10</sup> as  $GA(G) = \sum_{i=1}^{|E(G)|} \xi_i$  in which, for the edge  $e_i = u_i v_i \in E(G)$ ,

$$\xi_i = \frac{2\sqrt{du_i dv_i}}{du_i + dv_i} \text{ and } du \text{ denoted to the degree of vertex } u. \text{ In this paper we compute some results}$$

about this new topological index. Herein, our notation is standard and taken from the standard book of graph theory<sup>11-19</sup>.

---

\* Corresponding author: jalali6834@gmail.com)

**2. Main results and discussion**

Before going to calculate the GA index for  $TUC_4C_8(S)$  nanotube, we must compute this new index, for some well-known class of graphs.

**Example 1.** Let  $C_n$  be a cycle on  $n$  vertices. We know all of vertices are of degree 2 and so,  $GA(G) = \sum_{e \in E} \frac{2\sqrt{2 \times 2}}{2+2} = n$ .

**Example 2.** Let  $S_n$  be a star on  $n + 1$  vertices (figure 1). One can see there are  $n$  vertices of degree 1 and a vertex of degree  $n$ . So,  $GA(G) = \sum_{e \in E} \frac{2\sqrt{du dv}}{du + dv} = \frac{2n\sqrt{n}}{n+1}$ .




Fig. 1. Star graph with  $n + 1$  vertices.

**Example 3.** Let  $GP(n, k)$  be generalized Petersen graph with parameters  $n$  and  $k$ , vertex set  $V = \{x_1, \dots, x_n, y_1, \dots, y_n\}$  and edge set

$E = \{x_1x_2, x_2x_3, \dots, x_nx_1, x_1y_1, x_2y_2, \dots, x_ny_n, y_1y_{k+1}, y_2y_{k+2}, \dots, y_ny_{k+n}\} \pmod n$  respectively. It

is easy to see that  $|E(GP(n, k))| = \begin{cases} 3n & \text{if } n \neq 2k \\ \frac{5n}{2} & \text{if } n = 2k \end{cases}$  and so, we have

$$GA(GP(n, k)) = \begin{cases} 3n & \text{if } n \neq 2k \\ \frac{3n}{2} + \frac{2n\sqrt{6}}{5} & \text{if } n = 2k \end{cases}$$




Fig. 2. Graph of wheel on  $n + 1$  vertices.

Now we compute the GA index of a  $TUC_4C_8(S)$  nanotube as described above. The GA index of the 2-dimensional lattice of  $TUC_4C_8(S)$  graph  $K = KTUC[p,q]$  (Figure 3) is also computed. Following Diudea<sup>20-24</sup>, we denote a  $TUC_4C_8(S)$  nanotube by  $G = GTUC[p,q]$ ,  $TUC_4C_8(R)$  nanotorus by  $H = HTUC[p,q]$  (Figures 4 and 5). It is easy to see that  $|V(K)| = |V(G)| = |V(H)| = 8pq$ ,  $|E(K)| = 12pq - 2p - 2q$ ,  $|E(G)| = 12pq - 2p$  and  $|E(H)| = 12pq$ . We begin with the molecular graph of  $K$  (Figure 3). One can see that there are three separate cases and the number of edges is different. Suppose  $e_1$ ,  $e_2$  and  $e_3$  are representative edges for these cases. We can see that  $\xi_1 = \xi_3 = 1$  and  $\xi_2 = \frac{2\sqrt{6}}{5}$ . By the definition of GA index and table 1 one can see that

$GA(K) = 12pq + (\frac{8\sqrt{6}}{5} - 6)(q - p) + 8 - \frac{16\sqrt{6}}{5}$ . We now consider the molecular graph  $G$  of Figure 4. This figure shows that there are two different cases and the number of edges is different. Suppose  $e_1$  and  $e_2$  are representatives of the different cases. One can see that  $\xi_1 = 1$  and  $\xi_2 = \frac{2\sqrt{6}}{5}$ . On the other hand, there are  $2p$  and  $4p$  similar edges for each of edges  $e_1$  and  $e_2$ , respectively. This implies that:

$$GA(G) = 12pq + (\frac{8\sqrt{6}}{5} - 6)p.$$

**Lemma 1.** For an arbitrary graph  $G$ ,  $GA(G) = |E(G)|$  if and only if  $G$  be a  $k$ -regular graph.

**proof.** If  $G$  be  $k$ -regular then it is easy to see that for every  $e \in V(G)$ ,  $\xi = 1$  and then  $GA(G) = |E(G)|$ . Conversely, suppose  $GA(G) = |E(G)|$ . So,  $\xi_1 + \xi_2 + \dots + \xi_{|E(G)|} = |E(G)|$ . This implies  $\xi_i = 1$  ( $1 \leq i \leq |E(G)|$ ) and proof is completed.

Now consider the Figure 5. Because this graph is 3-regular, by using lemma 1 we have:

$$GA(H) = |E(H)| = 12pq.$$

Table 1. Computing the  $\xi_i$  for the 2-dimensional lattice of  $TUC_4C_8(S)$  graph  $K = KTUC[p,q]$ .

No.	$\xi_i$	Type of Edges
$2p+2q+4$	1	$e_1$
$4(p+q-2)$	$\frac{2\sqrt{6}}{5}$	$e_2$
$12pq - 8p - 8q + 4$	1	$e_3$




Fig. 3. 2-Dimensional Lattice of  $TUC_4C_8(S)$  Nanotorus with  $p = 4$  and  $q = 2$ .




Fig. 4. The graph of  $TUC_4C_8(S)$  nanotube  $G = GTUC [p,q]$  with  $p = 4$  and  $q = 2$ .




Fig. 5. The 2-Dimensional Lattice of  $TUC_4C_8(S)$  Nanotorus.

## References

- [1] S. Iijima, Helical microtubules of graphitic carbon, *Nature*, **354**, 56 (1991).
- [2] Y. M. Yang and W. Y. Qiu, Molecular Design and Mathematical Analysis of Carbon Nanotube Links, *MATCH Commun. Math. Comput. Chem.*, **58**, 635 (2007).
- [3] A. T. Balaban, Carbon and its nets. Symmetry 2: unifying human understanding, Part 1., *Comput. Math. Appl.*, **17**, 397 (1989).
- [4] N. Trinajstić, *Chemical Graph Theory*, CRC Press, Boca Raton, FL., 1992.
- [5] I. Gutman, O. E. Polansky. *Mathematical Concepts in Organic Chemistry*, Springer-Verlag, New York, 1986.
- [6] M. A. Johnson, G. M. Maggiora, *Concepts and Applications of Molecular Similarity*, Wiley Interscience, New York, 1990.
- [7] P. E. John, A. E. Vizitiu, S. Cigher and M. V. Diudea, *MATCH Commun. Math. Comput. Chem.*, **57**, 479 (2007).
- [8] H. Wiener, *J. Am. Chem. Soc.*, **69**, 17 (1947).
- [9] H. Hosoya, *Disc. Appl. Math.*, **19**, 239 (1988).
- [10] D. Vukičević and B. Furtula, *J. Math. Chem.*, 2009, (DOI 10.1007/s10910-009-9520-x).
- [11] A. R. Ashrafi and M. Ghorbani, *MATCH Commun. Math. Comput. Chem.*, **59**(3), 595 (2008).
- [12] A. R. Ashrafi, H. Saati and M. Ghorbani, *Digest Journal of Nanomaterials and Biostructures*, **3**(4), 227 (2008).
- [13] A. R. Ashrafi, M. Jalali, M. Ghorbani and M. V. Diudea, *MATCH Commun. Math. Comput. Chem.*, **60**(3), 905 (2008).
- [14] M. Ghorbani and A. R. Ashrafi, *J. Comput. Theor. Nanosci.* **3**, 803 (2006).
- [15] A. R. Ashrafi, M. Jalali, M. Ghorbani and M. V. Diudea, *MATCH Commun. Math. Comput. Chem.*, **60**(3), 905 (2008).

- [16] A. R. Ashrafi, M. Ghorbani, and M. Hemmasi, *Digest Journal of Nanomaterials and Biostructures*, **4**(3), 483 (2009).
- [17] M. Ghorbani and M. Jalali, *MATCH Commun. Math. Comput. Chem.*, **62**, 353 (2009).
- [18] A. R. Ashrafi, M. Ghorbani and M. Jalali, *Optoelectronics and Advanced Materials-Rapid Communications*, **3**(8), 823 (2009).
- [19] A. R. Ashrafi and M. Ghorbani, *Optoelectronics and Advanced Materials-Rapid Communications*, **3**(6), 596 (2009).
- [20] M. V. Diudea, S. Cigher, A. E. Vizitiu, O. Ursu and P. E. John, *Croat. Chem. Acta*, **79**, 445 (2006).
- [21] A. E. Vizitiu, S. Cigher, M. V. Diudea, M. S. Florescu, *MATCH Commun. Math. Comput. Chem.*, **57**, 457 (2007).
- [22] M. V. Diudea, *Carpath. J. Math.*, **22**, 43 (2006).
- [23] P. E. John, A. E. Vizitiu, S. Cigher and M. V. Diudea, *MATCH Commun. Math. Comput. Chem.*, **57**, 479 (2007).
- [24] M. V. Diudea, *Fullerenes, Nanotubes, and Carbon Nanostructures*, **10**, 273 (2002).

CONCLUSION Topological indices are designed basically by transforming a molecular graph into a number[6].We compute a new distance based hyper Zagreb index and augmented Zagreb index for HAC 5 C 6 C 7 [p,q] and TUC 4 C 6 C 8 [p,q] nanotubes. First Author N.K.Raut, Dept of Physics,Majalgaon. College,Majalgaon (Dist:Beed) India., Email: rautNK87@gmail.com Second Author S.N.Ipper, Dept of Chemistry,Majalgaon College,Majalgaon (Dist:Beed) India. Documents Similar To Computing some topological indices of Nanotubes. Carousel Previous Carousel Next. A topological photonic crystal design directly generates light that carries orbital angular momentum with high quantum numbers. The beam contains several different states at the same time, promising integrated and multiplexed light sources. Hall, E. H. On a new action of the magnet on electric currents. Am. J. Math. Nonlinear light generation in topological nanostructures. Nat. Nanotechnol. To compute and study topological indices of nanostructures are a respected problem in nanotechnology. Quantitative structure-property and structure-activity relationships of the carbon nanotubes compounds necessitate expressions for the molecular topological features of these compounds. Topological indices are vital devices for investigating chemical compounds to comprehend the fundamental topology of chemical structures. The GA index is a topological index was defined as, in which degree of a vertex  $u$  denoted by  $d(u)$ . Atom bond connectivity index is another topological index defined as  $ABC(G)=\sum_{uv \in E} d(u)+d(v)-2/d(u)d(v)$ . In this paper we compute these topological indices for two types of nanocones. View. Show abstract.